arXiv:cond-mat/0305393v1 [cond-mat.dis-nn] 16 May 2003

On the existence of a finite-temperature transition in the two-dimensional gauge glass

Helmut G. Katzgraber
Theoretische Physik, ETH Hönggerberg, CH-8093 Zürich, Switzerland

(Dated: February 2, 2008)

Results from Monte Carlo simulations of the two-dimensional gauge glass supporting a zero-temperature transition are presented. A finite-size scaling analysis of the correlation length shows that the system does not exhibit spin-glass order at finite temperatures. These results are compared to earlier claims of a finite-temperature transition.

PACS numbers: 75.50.Lk, 75.40.Mg, 05.50.+q

The existence of a finite-temperature spin-glass transition in the two-dimensional gauge glass has been a source of great controversy. While the existence of a transition in three dimensions is undisputed,[1] several different approaches[2,3,4,5,6,7,8,9,10] have been made to determine the spin-glass ordering temperature T_c in two dimensions with different results. As the gauge glass in two dimensions is a model commonly used to describe the vortex-glass transition in dirty high-temperature superconductor thin films,[11] it is important to settle the question of whether the system exhibits spin-glass order at finite temperatures or not. Experimental results on YBCO thin films[12] show no vortex-glass order, whereas bulk materials do exhibit a transition to a glassy phase. A brief summary of the different approaches used to determine T_c is presented, followed by results from a novel technique[13] that strongly supports the absence of a glassy phase at finite temperatures in the two-dimensional gauge glass.

Early work by Fisher *et al.*[2] who study the size-dependence of domain-wall energies with a zero-temperature domain-wall renormalization group method, shows evidence of a zero-temperature transition in the two-dimensional gauge glass. Gingras,[14] as well as Akino and Kosterlitz,[15] who also study the stiffness exponent of the model, find agreeing results.

In the aforementioned work,[2] Fisher *et al.* also analyze the scaling of Binder ratios[16] of the order parameter and obtain further evidence of a zero-temperature transition. These results are in agreement with Monte Carlo simulations by Reger and Young[3] who conclude that the transition temperature is not finite by studying the scaling of the currents induced by a twist in the boundary conditions. Katzgraber and Young[9] find evidence of a zero-temperature transition for the two-dimensional gauge-glass Hamiltonian with periodic boundary conditions by studying the scaling of the currents and the spin-glass susceptibility. They also conclude that one can scale data for the susceptibility well for any positive T_c, probably due to the extra fitting parameter η, the anomalous dimension of the spin-glass order parameter q, and show that studying the scaling of the spin-glass susceptibility alone is not enough to obtain evidence of a finite-temperature transition. This may explain why similar simulations by Choi and Park,[7] who study the susceptibility only, find a finite-temperature transition with $T_c = 0.22 \pm 0.02$.

Furthermore, Kim[8] has studied the gauge glass with fluctuating twist boundary conditions[17] by means of resistively shunted junction (RSJ) dynamics and finds a finite T_c in agreement with the work of Choi and Park,[7] and with early results by Li[18] (challenged later by Simkin[19]). Surprisingly, Granato,[6] who also uses RSJ dynamics but with periodic boundary conditions, obtains $T_c = 0$.

More recently Holme *et al.*[10] have studied the two-dimensional gauge glass by means of Monte Carlo simulations with fluctuating twist boundary conditions. By introducing an extra anomalous exponent b they are able to scale data for the currents and the helicity modulus and so obtain a finite transition temperature close to 0.20.

As there are diverging opinions on the spin-glass transition temperature of the two-dimensional gauge glass, the issue is reconsidered in the present work using a different approach. The scaling of the finite-system correlation lengths is studied, giving compelling evidence that $T_c \approx 0$, at least for the system sizes studied here. These results are supported by data for the spin-glass susceptibility.

The gauge-glass Hamiltonian,[20] which describes a disordered granular superconductor, is given by

$$\mathcal{H} = -J \sum_{\langle i,j \rangle} \cos(\phi_i - \phi_j - A_{ij}), \qquad (1)$$

where the sum ranges over nearest neighbors on a square lattice of size $N = L^2$ and ϕ_i represent the angles of the XY spins. The A_{ij} are quenched random variables uniformly distributed between $[0, 2\pi]$ with the constraint that $A_{ij} = -A_{ji}$. In this work $J = 1$ and periodic boundary conditions are applied.

Previously[9] we estimated the critical temperature of the two-dimensional gauge glass by studying the scaling of the currents[21] induced by a twist in the boundary conditions. This has an advantage over studying the crossing of Binder ratios[16] because the latter will not splay enough for XY systems[1] at $T < T_c$, making it almost impossible to accurately determine the crossing point. Studying the currents requires some care as well, as these have different finite-size scaling forms[9] depending on a finite or zero transition temperature. Finally, the currents show strong statistical fluctuations and so require a large amount of



FIG. 1: Correlation length ξ_L divided by L for different system sizes. At all temperatures the data decrease with increasing L. This indicates that if T_c is finite, it must be less than 0.13, the lowest temperature simulated. The shaded region represents the area around $T \approx 0.20$ where the claimed (Refs. 7,8,10) finite-temperature transition takes place. The inset shows a scaling plot of the data for ξ_L/L plotted against $L^{1/\nu}T$ for $T_c = 0$ and $1/\nu = 0.39$. One can see that the data scale very well, especially for low temperatures.

disorder realizations. Therefore in this work the scaling of the correlation lengths is studied as these show little statistical fluctuations, have a simple finite-size scaling form and splay out well for $T < T_c$. This method has been applied with success to Ising,[13] as well as vector spin glasses.[22] The spin-glass order parameter for the gauge glass is given by

$$q = \frac{1}{N}\sum_j^N \exp[i(\phi_j^\alpha - \phi_j^\beta)] \; , \qquad (2)$$

where "α" and "β" represent two replicas of the system with the *same* disorder. Equation (2) is generalized to finite wave vectors $\mathbf{k}$:

$$q(\mathbf{k}) = \frac{1}{N}\sum_j^N \exp[i(\phi_j^\alpha - \phi_j^\beta) + i\mathbf{k}\cdot\mathbf{R}_j] \; . \qquad (3)$$

The wave vector dependent spin-glass susceptibility $\chi_{\rm SG}(\mathbf{k})$ is then given by

$$\chi_{\rm SG}(\mathbf{k}) = N[\langle |q(\mathbf{k})|^2\rangle]_{\rm av} \; . \qquad (4)$$

Here, $\langle\cdots\rangle$ denotes a thermal and $[\cdots]_{\rm av}$ a disorder average. The finite-system correlation length ξ_L is determined from an expansion[13,23,24] of the spin-glass susceptibility for small $\mathbf{k}\xi_L$ in the framework of an Ornstein-Zernicke approximation[25]

$$[\chi_{\rm SG}(\mathbf{k})/\chi_{\rm SG}(\mathbf{0})]^{-1} = 1 + (\mathbf{k}\xi_L)^2 + O[(\mathbf{k}\xi_L)^4] \; . \qquad (5)$$

Keeping only the leading term in the expansion and taking into account the lattice periodicity as well as corrections to scaling, we obtain

$$\xi_L = \frac{1}{2\sin(|\mathbf{k}_{\rm min}|/2)}\left[\frac{\chi_{\rm SG}(\mathbf{0})}{\chi_{\rm SG}(\mathbf{k}_{\rm min})} - 1\right]^{1/2} \; , \qquad (6)$$

where $\mathbf{k}_{\rm min} = (2\pi/L, 0, 0)$ is the smallest nonzero wave vector. In Eq. (6), $\chi_{\rm SG}(\mathbf{0})$ is the standard spin-glass susceptibility.

The finite-size scaling of ξ_L can be understood by studying the finite-size scaling of the spin-glass susceptibility extended to finite wave vectors k

$$\chi_{\rm SG}(T, L, k) = L^{2-\eta}\tilde{C}[L^{1/\nu}(T - T_c), kL] \; . \qquad (7)$$

From Eq. (5) $k_{\rm min}^2\xi_L^2 \sim \chi_{\rm SG}(0)/\chi_{\rm SG}(k_{\rm min})$ up to an additive constant, and so the L-dependent prefactor in Eq. (7) cancels. The dependence of $\chi_{\rm SG}$ on kL drops out as well, as in Eq. (6) $k = k_{\rm min} = 2\pi/L$. We therefore obtain the simple finite-size scaling form

$$\xi_L/L = \tilde{X}[L^{1/\nu}(T - T_c)] \; , \qquad (8)$$

and so, the introduction of an anomalous exponent[10] is not plausible. This has also the advantage that the finite-size scaling has few parameters. From Eq. (8) we see that data for different system sizes L will cross at $T = T_c$. If $T_c = 0$, one expects that data for different L will decrease with increasing L and not cross for any finite temperature.

For the simulations the parallel tempering Monte Carlo method[26,27] is used. Because the equilibration test for short-range spin glasses introduced by Katzgraber *et al.*[22,28,29] does not work for the gauge glass[29] as the disorder is not Gaussian, equilibration is tested by the traditional technique of requiring that different observables are independent of the number of Monte Carlo steps. By doubling the number of Monte Carlo sweeps between each measurement until the last three agree within error bars, equilibration is ensured. The equilibration times used for the different system sizes are listed in Ref. 9, Table I. For the largest size studied, $L_{\rm max} = 16$, 10^3 disorder realizations are performed, whereas for the smaller sizes L, $10^3(L_{\rm max}/L)$ realizations. The lowest temperature studied is $T = 0.13$, well below the claimed[7,8,10] transition at $T \approx 0.20$. Here, 30 temperature replicas are used with the highest temperature being $T = 1.058$.

Figure 1 shows data for ξ_L/L for different system sizes L. The data decrease with increasing L for all temperatures. In particular, there is *no* crossing of the data around $T \approx 0.20$ (shaded area) where the spin-glass transition is supposed to take place.[7,8,10] The inset of Fig. 1 shows a finite-size scaling plot of the data according to Eq. (8). One can see that the data scale very well for $T_c = 0$ and $1/\nu = 0.39 \pm 0.02$, values found in Ref. 9 by scaling the currents and the spin-glass susceptibility. The error bar in $1/\nu$ is determined by varying its value until the data do not scale well. An attempt to scale the



FIG. 2: Data for $\ln(\chi_{SG}/L^2)$ vs $\ln(L)$ for different temperatures. At T_c one expects $\chi_{SG} \sim L^{2-\eta}$. The data show a strong downward curvature indicating that $T > T_c$ for the range of temperatures shown. This is emphasized in the inset where the curvature (from a second-order polynomial fit) is displayed. At T_c one expects the curvature to cross zero. Here the data extrapolate to $T_c \approx 0$ (the dashed line represents a second-order polynomial fit extrapolation). The shaded area shows the region where one would expect the curvature to cross zero if $T_c \approx 0.2$. A finite-size scaling analysis of the spin-glass susceptibility according to Eq. (9) works well for low enough temperatures with $T_c = 0$, $\eta = 0$, and $1/\nu = 0.39$, in agreement with results from Ref. 9.

data in Fig. 1 to any value of a finite T_c, in particular 0.20, fails for any choice of ν.

Standard finite-size scaling predicts that the spin-glass susceptibility scales as[30,31]

$$\chi_{SG} = L^{2-\eta}\tilde{C}[L^{1/\nu}(T - T_c)] \; , \qquad (9)$$

which means that at $T = T_c$ it should scale as a power law $\sim L^{2-\eta}$. In Fig. 2 data for $\ln(\chi_{SG}/L^2)$ vs $\ln(L)$ is shown. If the data were at T_c, one would expect a straight line. In particular, $T = 0.20$ shows a strong downward curvature indicating that the system is *above* the critical transition temperature. A fit of the data in Fig. 2 to a straight line for the different temperatures yields quality of fit probabilities[32] between 10^{-9} for the lowest, and 10^{-20} for the highest temperature, respectively. A second-order polynomial fit to the data, instead, has quality of fit probabilities ~ 0.9. The curvature of the second-order fits is determined and shown in the inset of Fig. 2. Because at $T < T_c$ $[T > T_c]$ one expects $\chi_{SG}(L)$ to bend upward [downward], the curvature has to go through zero at T_c. The data in the inset of Fig. 2 do not cross the horizontal axis for the temperature range studied and extrapolate within error bars to $T_c \approx 0$. This is in agreement with results by Katzgraber and Young[9] who obtain good scaling of the susceptibility at low temperatures for $T_c = 0$, $\eta = 0$, and $1/\nu = 0.39$.

To conclude, evidence of a zero-temperature transition in the two-dimensional gauge glass with periodic boundary conditions is presented by studying the finite-size scaling of the correlation lengths. This approach is better than previous methods used because the correlation lengths show small statistical fluctuations and have a simple finite-size scaling form. In addition, the curvature of the spin-glass susceptibility as a function of system size indicates that there is no spin-glass transition at $T \approx 0.20$, at least for the system sizes studied. Moreover the curvature extrapolates within error bars to $T_c \approx 0$. These results are in agreement with experimental data on YBCO thin films,[12] where a finite-T is absent, and different numerical approaches.[2,3,9,15] In particular, it is shown that the system does *not* exhibit a spin-glass transition at $T \approx 0.20$, as claimed in Refs. 7, 8, and 10. It is noteworthy that Refs. 8 and 10 find the aforementioned finite temperature transition by studying the gauge glass with fluctuating twist boundary conditions instead of the commonly used periodic boundary conditions. Could this be one source of the discrepancies?

Finite-size effects can influence the data, and so an analysis with larger system sizes is desirable. Nevertheless, the data presented here strongly support the absence of a transition at $T \approx 0.20$, although a transition at $T < 0.13$, the lowest temperature studied, while unlikely the case, cannot be ruled out completely.

I would like to thank G. Blatter, I. Campbell, M. Troyer, and A. P. Young for helpful discussions and to K. Tran for carefully reading the manuscript. The simulations were performed on the Asgard cluster at ETH Zürich.

---

[1] T. Olson and A. P. Young, *Finite temperature ordering in the three-dimensional gauge glass*, Phys. Rev. B **61**, 12467 (2000), (cond-mat/991229).

[2] M. P. A. Fisher, T. A. Tokuyasu, and A. P. Young, *Vortex variable-range-hopping resistivity in superconducting films*, Phys. Rev. Lett. **66**, 2931 (1991).

[3] J. D. Reger and A. P. Young, *Monte Carlo study of a vortex glass model*, J. Phys. A **26**, 1067 (1993).

[4] R. A. Hyman, M. Wallin, M. P. A. Fisher, S. M. Girvin, and A. P. Young, *Current-voltage characteristics of two-dimensional vortex-glass models*, Phys. Rev. B **51**, 15304 (1995).

[5] J. M. Kosterlitz and M. V. Simkin, *Numerical Study of a Superconducting Glass Model*, Phys. Rev. Lett. **79**, 1098

(1997).

[6] E. Granato, *Current-voltage scaling of chiral and gauge-glass models of two-dimensional superconductors*, Phys. Rev. B **58**, 11161 (1998).

[7] M. Y. Choi and S. Y. Park, *Phase transition in the two-dimensional gauge glass*, Phys. Rev. B **60**, 4070 (1999).

[8] B. J. Kim, *Finite-temperature resistive transition in the two-dimensional XY gauge glass model*, Phys. Rev. B **62**, 644 (2000).

[9] H. G. Katzgraber and A. P. Young, *Numerical studies of the two- and three-dimensional gauge glass at low temperature*, Phys. Rev. B **66**, 224507 (2002), (cond-mat/0205206).

[10] P. Holme, B. J. Kim, and P. Minnhagen, *Phase transitions in the two-dimensional random gauge XY model*, Phys. Rev. B **67**, 104570 (2003), (cond-mat/0301279).

[11] G. Blatter, M. V. Feigelman, V. B. Geshkenbein, A. I. Larkin, and V. M. Vinokur, *Vortices in high-temperature superconductors*, Rev. Mod. Phys. **66**, 1125 (1994).

[12] C. Dekker, P. M. J. Wöltgens, R. H. Koch, B. W. Hussey, and A. Gupta, *Absence of a Finite-Temperature Vortex-Glass Phase Transition in Two-Dimensional $YBa_2Cu_3O_{7-\delta}$ Films*, Phys. Rev. Lett. **69**, 2717 (1992).

[13] H. G. Ballesteros, A. Cruz, L. Fernandez, V. Martin-Mayor, J. Pech, J. J. Ruiz-Lorenzo, A. Tarancon, P. Tellez, C. L. Ullod, and C. Ungil, *Critical behavior of the three-dimensional Ising spin glass*, Phys. Rev. B **62**, 14237 (2000), (cond-mat/0006211).

[14] M. J. P. Gingras, *Numerical study of vortex-glass order in random-superconductor and related spin-glass models*, Phys. Rev. B **45**, 7547 (1992).

[15] N. Akino and J. M. Kosterlitz, *Domain wall renormalization group study of xy model with quenched random phase shifts*, Phys. Rev. B **66**, 054536 (2002), (cond-mat/0203299).

[16] K. Binder, *Critical properties from Monte Carlo coarse graining and renormalization*, Phys. Rev. Lett. **47**, 693 (1981).

[17] P. Olsson, *Effective vortex interaction in the two-dimensional XY model*, Phys. Rev. B **46**, 14598 (1992).

[18] Y. Li, *Voltage-Current Characteristics of the Two-Dimensional Gauge Glass Model*, Phys. Rev. Lett. **69**, 1819 (1992).

[19] M. V. Simkin, *Comment on "Voltage-Current Characteristics of the Two-Dimensional Gauge Glass Model"* (1996), (cond-mat/9604178).

[20] W. Y. Shih, C. Ebner, and D. Stroud, *Frustration and disorder in granular superconductors*, Phys. Rev. B **30**, 1134 (1984).

[21] J. D. Reger, T. A. Tokuyasu, A. P. Young, and M. P. A. Fisher, *Vortex-glass transition in three dimensions*, Phys. Rev. B **44**, 7147 (1991).

[22] L. W. Lee and A. P. Young, *Single spin- and chiral-glass transition in vector spin glasses in three dimensions* (2003), (cond-mat/0302371).

[23] F. Cooper, B. Freedman, and D. Preston, *Solving $\phi^4_{1,2}$ theory with Monte Carlo*, Nucl. Phys. B **210**, 210 (1982).

[24] The expression for the finite-system correlation length also follows from the Ornstein-Zernicke equation for the two-point function (spin-glass susceptibility).

[25] V. Martín-Mayor, A. Pelissetto, and E. Vicari, *Critical structure factor in Ising systems*, Phys. Rev. E **66**, 26112 (2002).

[26] K. Hukushima and K. Nemoto, *Exchange Monte Carlo method and application to spin glass simulations*, J. Phys. Soc. Jpn. **65**, 1604 (1996).

[27] E. Marinari, *Optimized Monte Carlo methods*, in *Advances in Computer Simulation*, edited by J. Kertész and I. Kondor (Springer-Verlag, Berlin, 1998), p. 50, (cond-mat/9612010).

[28] H. G. Katzgraber, M. Palassini, and A. P. Young, *Monte Carlo simulations of spin glasses at low temperatures*, Phys. Rev. B **63**, 184422 (2001), (cond-mat/0007113).

[29] H. G. Katzgraber and A. P. Young, *Monte Carlo simulations of the four dimensional XY spin-glass at low temperatures*, Phys. Rev. B **65**, 214401 (2002), (cond-mat/0108320).

[30] In Ref. 9 a finite-size scaling plot of the spin-glass susceptibility is performed with $1/\nu = 0.39$, $T_c = 0$, and $\eta = 0$. The data scale well at the lowest temperatures studied. For higher temperatures, strong corrections to scaling are visible. In fact, the best fit "by eye" yields $1/\nu = 0.50$. This slight discrepancy can be explained by recent work (Ref. 31) of Carter *et al.* who show that for a zero-temperature transition the scaling can have considerable corrections due to large corrections to scaling in the expression for the correlation length.

[31] A. C. Carter, A. J. Bray, and M. A. Moore, *On the use of finite-size scaling to measure spin-glass exponents* (2003), (cond-mat/0302207).

[32] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical Recipes in C* (Cambridge University Press, Cambridge, 1995).